

Manual

CircLigase II ssDNA Ligase

For Research Use Only. Not for use in diagnostic procedures.

CircLigase™ II ssDNA Ligase is part of the Epicentre™ product line, known for its unique genomics kits, enzymes, and reagents which offer high quality and reliable performance.

Manual

CircLigase II ssDNA Ligase

Contents

1. Introduction	3
2. Product designations and kit components	3
3. Product specifications	3
4. Applications	4
5. General considerations	4
6. Kit procedure	5
7. Further support	6

Manual

CircLigase II ssDNA Ligase

1. Introduction

[CircLigase II ssDNA Ligase](#)* is a thermostable ligase that catalyses intramolecular ligation (i.e., circularisation) of single-stranded DNA (ssDNA) and single-stranded RNA (ssRNA) substrates that have both a 5'-monophosphate and a 3'-hydroxyl group. Linear ssDNAs and ssRNAs of greater than 30-35 bases (sequence dependent) are circularised by CircLigase ssDNA Ligase. Under standard reaction conditions, virtually no linear concatamers or circular concatamers are produced.

2. Product designations and kit components

Product	Kit size	Catalogue number	Reagent description	Part number	Volume
CircLigase II ssDNA Ligase	1,000 units	CL9021K	CircLigase II ssDNA Ligase (100 U/ μ L)	E0129-100D3	10 μ L
			Betaine (5M)	SS000026-D4	50 μ L
			MnCl ₂ (50 mM)	SS000578-D2	20 μ L
			CircLigase ssDNA Control (2 pmole/ μ L)	SS000592-D1	10 μ L
			Nuclease-Free Water, Sterile	SS000772-D3	1 mL
			CircLigase II 10X Reaction Buffer	SS000881-D1	50 μ L
	5,000 units	CL9025K	CircLigase II ssDNA Ligase (100 U/ μ L)	E0129-100D4	50 μ L
			Betaine (5M)	SS000026-D5	250 μ L
			MnCl ₂ (50 mM)	SS000578-D3	75 μ L
			CircLigase ssDNA Control (2 pmole/ μ L)	SS000592-D2	25 μ L
			Nuclease-Free Water, Sterile	SS000772-D3	1 mL
			CircLigase II 10X Reaction Buffer	SS000881-D2	150 μ L

3. Product specifications

Storage: Store only at -20 °C in a freezer without a defrost cycle.

Storage buffer: CircLigase II ssDNA Ligase is supplied in a 50% glycerol solution containing 50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA, 1 mM dithiothreitol (DTT) and 0.1% Triton X-100 (Rohm & Haas).

Unit definition: One unit of CircLigase II ssDNA Ligase converts 1 pmol of a linear 5'-phosphorylated CircLigase II Control Oligo (55 mer) into an exonuclease I-resistant circular ssDNA in 1 hour at 60 °C under standard assay conditions.

CircLigase II 10X Reaction Buffer: 0.33 M Tris-acetate (pH 7.5), 0.66 M potassium acetate and 5 mM DTT.

For circularisation of ssDNA, we recommend adding MnCl₂ to a final concentration of 2.5 mM.

Contaminating activity assays: CircLigase II ssDNA Ligase is free of detectable DNA exonuclease and endonuclease, and RNase activities.

Manual

CircLigase II ssDNA Ligase

4. Applications

- Production of single-stranded DNA templates for rolling-circle replication or rolling-circle transcription experiments and next-generation sequencing.
- Production of circular ssRNA >30-35 nt (sequence dependent).

5. General considerations

1. **Substrate requirements:** The circularisation reaction requires ssDNA or ssRNA with 5'-phosphate and 3'-hydroxyl groups. The standard CircLigase II reaction uses 10 pmol of linear ssDNA.
2. **Substrate size:** The ssDNA or ssRNA must be at least ~15 bases in length. Substrates such as single-stranded oligodeoxynucleotides and single-stranded cDNAs can be ligated by the enzyme.
3. **Manganese:** For circularisation of ssDNA or ssRNA, such as oligodeoxynucleotides or cDNA, add MnCl_2 to a final concentration of 2.5 mM. A tube of MnCl_2 is included.
4. **Magnesium:** In general, circularisation is better in the absence of magnesium.
5. **Amount of CircLigase II ssDNA Ligase in the reaction:** The standard reaction conditions (section 6.1) use 100 U of the CircLigase II enzyme per 20 μL reaction (~1 μM enzyme and 0.5 μM ssDNA substrate). For custom ligation reactions, we recommend maintaining the enzyme concentration in excess of the substrate concentration.
6. **Sequence dependence:** Our results indicate that the sequence of the ssDNA can strongly influence the efficiency of the circularisation reaction.
7. **Reaction time:** The CircLigase II ssDNA circularisation reaction is typically complete in 60 minutes. However, increasing the reaction time may improve the yield of circular DNA with difficult-to-ligate ssDNA substrates. In some cases, the molar concentration of the substrate DNA may be high enough that ligation may not be complete; it may be helpful to add 1 μL of 1 mM ATP to allow the ligation reaction to proceed to completion.
8. **Betaine:** Betaine is not necessary for circularisation of easy-to-ligate ssDNA molecules. However, we have found that difficult-to-ligate ssDNA substrates can be circularised by including betaine at a final concentration of 1 M in the ligation reaction. A separate tube of betaine is provided in the kit to enable optimisation of the betaine concentration, if necessary.
9. **Difficult substrates:** Some ssDNAs or ssRNAs are inefficiently circularised in the standard reaction (section 6.1). The yield of circular ssDNA from a difficult-to-ligate substrate may be increased by increasing the concentration of CircLigase II ssDNA Ligase in the reaction, lengthening the reaction time (see note 7, above), or by adding betaine to the reaction (see note 8, above).
10. **The CircLigase II ssDNA Control Oligo:** The CircLigase II ssDNA Control Oligo provided in the kit is a 55 base oligodeoxynucleotide containing both 5'-phosphate and 3'-hydroxyl ends. Under standard reaction conditions (10 pmole Control Oligo, 100 U CircLigase II ssDNA Ligase, 2.5 mM MnCl_2 , 1 hour reaction), the linear Control Oligo is converted to circular ssDNA.

Manual

CircLigase II ssDNA Ligase

6. Kit procedure

6.1. Ligation reaction

1. Combine the following reaction components:

	Final Concentration
x μL Nuclease-Free Water	—
10 pmol Single-stranded DNA or RNA template	0.5 pmol/ μL
2 μL CircLigase II 10X Reaction Buffer	1X
1 μL 50 mM MnCl_2	2.5 mM
4 μL 5 M Betaine (optional)	1 M
1 μL CircLigase II ssDNA Ligase (100 U)	5 U/ μL
<hr/>	
20 μL Total reaction volume	

2. Incubate the reaction at 60 °C for 1 hour.

NOTE: Longer incubation times may improve the yield of circular ssDNA for difficult-to-ligate ssDNAs. For example, we have observed that the ligation reaction with some ssDNAs went to completion in the presence of 1 M betaine after 16 hours of incubation.

3. Incubate the reaction at 80 °C for 10 minutes to inactivate the CircLigase II ssDNA Ligase.

6.2. Gel analysis of the ligation reaction

The efficiency of a CircLigase II ligation reaction can be readily assessed by gel electrophoresis. When ligating oligos, load approximately 1 pmol of linear ssDNA substrate in one gel lane and 2 μL of the standard CircLigase II reaction mixture into an adjacent gel lane of a **20% acrylamide/8 M urea denaturing gel**. Run the gel and stain with an appropriate DNA-binding dye. The circularised ssDNA product migrates slower, above, the linear ssDNA band (see figure 1). In some instances, the adenylated oligo-intermediate can be seen as a band just above the linear ssDNA.




Figure 1. CircLigase II ssDNA Ligase converts linear ssDNA into closed circular ssDNA. A 71-nucleotide ssDNA oligo was converted to a circular ssDNA. Lane 1, DNA markers; lane 2, 71-nucleotide linear ssDNA oligo; lane 3, circularisation proceeds through an adenylated intermediate; lane 4, closed-circular ssDNA reaction product.

Manual

CircLigase II ssDNA Ligase

6.3. Removing the linear ssDNA template and adenylated intermediate from the reaction

Once the CircLigase II reaction has been terminated, the remaining linear ssDNA substrate and linear single-stranded adenylated intermediate can be removed by treatment with Exonuclease I (which digests linear ssDNA) and Exonuclease III (which digests linear double-stranded DNA). The circular ssDNA is resistant to these exonucleases, while the linear ssDNA and adenylated intermediate are digested. Single-stranded linear nucleic acids that were not circularised in the CircLigase reaction can be removed by digestion with Exonuclease I (for DNA), or Terminator™ Exonuclease or RNase R (for RNA).

Most linear ssDNA and adenylated-intermediate can be eliminated by addition of 20 U of Exonuclease I, followed by incubation at 37 °C for 45 minutes.

However, if the linear ssDNA substrate contains hairpins or other secondary structures, treatment with both Exonuclease I and Exonuclease III is recommended. We suggest incubating a standard ligation reaction mixture with 10 U of Exonuclease I and 100 U of Exonuclease III at 37 °C for 45 minutes.

7. Further support

If you require any further support, please do not hesitate to contact our Technical Support Team: techsupport@lgcgroup.com.

*CircLigase II ssDNA Ligase is covered by intellectual property rights licensed to LGC Biosearch Technologies. The purchase of this product conveys to the buyer the nontransferable right to use the purchased product and components of the product in research conducted by the buyer. The buyer cannot sell or otherwise transfer this product or its components to a third party and in particular, no rights are conveyed to the buyer to use the product or its components for commercial use purpose other than for research to gain information that is used by the buyer.



X **f in** @LGCBiosearch | biosearchtech.com

All trademarks and registered trademarks mentioned herein are the property of their respective owners. All other trademarks and registered trademarks are the property of LGC and its subsidiaries. Specifications, terms and pricing are subject to change. Not all products are available in all countries. Please consult your local sales representative for details. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or any retrieval system, without the written permission of the copyright holder. © LGC Limited, 2024. All rights reserved. GEN/939/EK/0924

For Research Use Only. Not for use in diagnostic procedures.



BIOSEARCH™
TECHNOLOGIES

GENOMIC ANALYSIS BY LGC